

Ranunculus jiguanshanicus (Ranunculaceae), a new species from Sichuan, China

Wen-Qun Fei^{1,2}, Qiong Yuan^{1,3}, Qin-Er Yang^{1,3}

1 Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510655, Guangdong, China **2** University of Chinese Academy of Sciences, Beijing 100049, China **3** Center of Conservation Biology, Core Botanical Gardens, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510655, Guangdong, China

Corresponding author: Qiong Yuan (yuanqiong@scib.ac.cn)

Academic editor: Marco Pellegrini | Received 13 October 2022 | Accepted 6 January 2023 | Published 25 January 2023

Citation: Fei W-Q, Yuan Q, Yang Q-E (2023) *Ranunculus jiguanshanicus* (Ranunculaceae), a new species from Sichuan, China. PhytoKeys 219: 57–75. <https://doi.org/10.3897/phytokeys.219.96266>

Abstract

Ranunculus jiguanshanicus (Ranunculaceae), a new species from Chongzhou in Sichuan province, China, is here described and illustrated. The new species is easily distinguishable from other Chinese members of the genus by an array of characters, including small stature, glabrous and prostrate stems, 3-foliolate leaves with obvious petiolules (3–5 mm long), unequally 3-sected leaflets, lanceolate to linear ultimate leaflet segments, small flowers (5.2–6 mm in diameter), and long styles in the carpels and achenes (ca. 0.8 mm long). A distribution map of this new species is also provided.

Keywords

Asia, buttercups, Ranunculales, *Ranunculus glareosus*, *Ranunculus pegaus*

Introduction

Ranunculus L., comprising approximately 600 species, is the largest genus in the Ranunculaceae and is widely distributed in all continents (Tamura 1995; Hörandl et al. 2005; Paun et al. 2005; Hörandl and Emadzade 2012). In China, one of the centers of species diversity of *Ranunculus*, more than 150 species and 30 varieties are currently recognized in the genus (Wang 1995a, b, 1996, 2007, 2008, 2013, 2015, 2016, 2018, 2019a, b, 2022; Yang 2000; Wang and Gilbert 2001; Wang and Liao 2009; Luo and Zhao 2013; Wang and Chen 2015; Wang et al. 2016; Yuan and Yang 2017a, b, c; Zhang et al. 2020; Fei et al. 2022, 2023a, b).

During a survey of herbarium specimens of *Ranunculus* from China for the first author's Ph.D. dissertation project, one gathering, W.B. Ju, L. Zhang & D.K. Chen AZH01290 (CDBI) (Fig. 1), from Jiguan Shan in the Anzihe Nature Reserve in Chongzhou, Sichuan province, China, caught our attention. This gathering had been previously identified on the determination slips as *R. glareosus* Hand.-Mazz., a species distributed in China's Qinghai, Sichuan, Xizang (Tibet) and Yunnan (Handel-Mazzetti 1931; Liou 1980; Wang and Gilbert 2001). The plants on the two sheets of the gathering, which have unique leaf morphology and very small flowers, certainly do not belong to *R. glareosus* or any other members of *Ranunculus* currently known for China.

During a botanical expedition to Sichuan from June to July 2022, we successfully discovered a flowering population of this species in early June on Jiguan Shan in Chongzhou, where the gathering W.B. Ju, L. Zhang & D.K. Chen AZH01290 was made. Moreover, we discovered a fruiting population in early July on Xiling Xue Shan in Dayi, a mountain closely adjacent to Jiguan Shan. Based on our observations of living plants in the wild, we confirmed all the diagnostic characters of the new species observed from the herbarium specimens and determined that the gathering and the two populations in question represent a new species. Morphologically, this new species is somewhat similar to *R. pegaeus* Hand.-Mazz., a species occurring in southwestern China (Sichuan, Xizang, Yunnan), India (Sikkim) and Nepal (Handel-Mazzetti 1939; Liou 1980; Wang and Gilbert 2001), but differs by an array of characters. It is described below as *R. jiguanshanicus*.

Materials and methods

For morphological comparison, we critically examined specimens or high-resolution specimen images of *Ranunculus glareosus*, *R. jiguanshanicus* and *R. pegaeus* at CDBI, E, KUN, PE, and WU (acronyms according to Thiers 2022). We also observed living plants in one population each in *R. glareosus* (Menyuan in Qinghai province) and *R. pegaeus* (Maoxian in Sichuan province) and those in two populations of *R. jiguanshanicus* (Chongzhou and Dayi in Sichuan province). The morphological description of *R. jiguanshanicus* was based on observations of both herbarium specimens and living plants in the wild.

Taxonomy

Ranunculus jiguanshanicus W.Q.Fei, Q.Yuan & Q.E.Yang, sp. nov.

urn:lsid:ipni.org:names:77312739-1

Figs 1–5

Diagnosis. *Ranunculus jiguanshanicus* is readily distinguishable from all other Chinese species of *Ranunculus* by a unique array of characters, including small stature, glabrous and prostrate stems, 3-foliolate leaves with obvious petiolules (3–5 mm long), unequally 3-sected leaflets, lanceolate to linear ultimate leaflet segments, small flowers (5.2–6 mm in diameter), and long styles in the carpels and achenes (ca. 0.8 mm long).

Type. CHINA. Sichuan province: Chongzhou, Anzihe Nature Reserve, Jiguan Shan, 30°46'5.8"N, 103°10'21.93"E, alt. 2998 m, among moss on rocks or rocky cliffs in moist places in fir forests, 10 June 2022, W.Q. Fei 581 (holotype: IBSC; isotypes: IBSC, PE).

Description. *Herbs* perennial, terrestrial or rupicolous. *Roots* 2–5, 6–10 cm long, fibrous, slender, slightly thickened at base. *Stems* 7–15 cm long, prostrate, glabrous, unbranched to few-branched. *Basal leaves* 2–5, 3-foliolate, long-petiolate; petioles 2–4 cm long, glabrous; blades 0.8–1 × 0.8–1.3 cm, suborbicular, thinly chartaceous, adaxially green, abaxially light green, both sides glabrous; leaflets 3, unequally 3-sected,



Figure 1. Two specimens of *Ranunculus jiguanshanicus* sp. nov. (**A**, **B**) previously misidentified as *R. glareosus*. China, Sichuan province, Chongzhou, Anzihe Nature Reserve, Jiguan Shan, 30°46'9.75"N, 103°10'11.65"E, alt. 2939 m, on rocks in fir forest, 16 June 2016, W.B. Ju, L. Zhang & D.K. Chen AZH01290 (CDBI). Insets: leaf blades.



Figure 2. *Ranunculus jiguanshanicus* sp. nov. in the wild **A, B** habitat **C** habit. The left plant (at fruiting stage) in **B** photographed by De-Chang Meng from Xiling Xue Shan in Dayi, Sichuan province, and the right two plants (at flowering stage) photographed by Wen-Qun Fei from Jiguan Shan in the Anzihe Nature Reserve in Chongzhou, Sichuan province.

petiolulate, petiolules 3–5 mm long, ultimate leaflet segments 3–4 × 0.8–1.2 mm, narrowly lanceolate to linear, margin entire, apex 1–2-denticulate to 1–2-cleft. **Lower cauline leaves** 2–3, similar to basal ones but smaller. **Upper cauline leaves** 1–2, 3-foliolate, subsessile or sessile, adaxially glabrous or sparsely puberulous, abaxially glabrous, central leaflet 4.5–5 × 1–1.2 mm, narrowly lanceolate to linear, margin entire, lateral leaflets




Figure 3. *Ranunculus jiguanshanicus* sp. nov. in the wild **A** roots **B** portion of stem **C** leaves **D** flower (lateral view) **E** flower (top view) **F** sepal (left: abaxial side; right: adaxial side) **G** petal (left: adaxial side; right: abaxial side) **H** stamens **I** carpels **J** aggregate fruit **K** achenes **L** receptacle. **A–H** photographed by Wen-Qun Fei from the population on Jiguan Shan in Chongzhou, Sichuan province and **I–L** photographed by De-Chang Meng from the population on Xiling Xue Shan in Dayi, Sichuan province.

entire, 1–2-lobate or 2–3-sected, ultimate leaflet segments $3\text{--}3.5 \times 1\text{--}1.2$ mm, narrowly lanceolate to linear. **Inflorescences** terminal, 1(–2)-flowered. **Flowers** 5.2–6 mm in diameter; pedicels 1–2 cm long, glabrous or sparsely puberulous; receptacles ca. 1.2 mm long, clavate, glabrous; sepals 5, $2.2\text{--}2.5 \times 1.5\text{--}1.8$ mm, elliptic to obovate, patent, green tinged with yellowish, concave, both sides glabrous; petals 5(–6), $3.2\text{--}3.5 \times 1.8\text{--}2$ mm, obovate, yellow, glabrous, apex rounded, nectary pit without




Figure 4. Holotype sheet of *Ranunculus jiguanshanicus* sp. nov.

a scale, claws ca. 0.4 mm long; stamens 6–8, ca. 2 mm long, filaments ca. 1.5 mm long, narrowly linear, anthers ca. 0.5 mm long, oblong; gynoecium subglobose; carpels 8–12, ovaries ca. 0.8 × 0.6 mm, ovoid, laterally flattened, biconvex, glabrous, styles ca. 0.8 mm long, glabrous, apex recurved. **Aggregate fruit** ca. 4 × 4.2 mm, subglobose; achenes ca. 1.2 × 1 mm, widely ovoid, laterally flattened, biconvex, glabrous, styles ca. 0.8 mm long, persistent, glabrous, apex recurved.

Etymology. The specific epithet refers to the type locality of the new species, i.e. Jiguan Shan in the Anzihe Nature Reserve in Chongzhou, Sichuan province, China.




Figure 5. Isotype (A–C) and paratype (D) sheets of *Ranunculus jiguanshanicus* sp. nov.

Phenology. Flowering in early June; fruiting at the end of June.

Distribution and habitat. *Ranunculus jiguanshanicus* is currently known from its type locality, i.e., Jiguan Shan in the Anzihe Nature Reserve in Chongzhou, and from the closely adjacent Xiling Xue Shan in Dayi, both in Sichuan province, China (Fig. 6).




Figure 6. Distribution of *Ranunculus jiguanishanicus* sp. nov. (black square). Arrow indicates the type locality.



Figure 7. Type sheets (A, B) of *Ranunculus pegasus*.



Figure 8. *Ranunculus pegaeus* in the wild (China, Sichuan, Maoxian) **A, B** habitat **C** habit. Photographed by Wen-Qun Fei.

It grows among moss on rocks or rocky cliffs in moist places in fir forests at altitudes of 2900–3150 m above sea level.

Conservation status. *Ranunculus jiguanshanicus* is currently known only from two populations in Sichuan province, China. The Chongzhou population consists of ca. 150 individuals within an area of less than 10 m². The size of the Dayi population remains unknown. The conservation status of *R. jiguanshanicus* should better be categorized as “Data Deficient (DD)” before adequate information of this species is acquired (IUCN Standards and Petitions Committee 2022).

Discussion. *Ranunculus jiguanishanicus* is readily assigned to *R.* sect. *Ranunculus* due to its swollen achenes with a distinct beak and receptacles hardly enlarged after anthesis. In his infrageneric classification of the Chinese *Ranunculus*, Wang (1995a, b) placed almost all the alpine species within this section under the name *R.* sect. *Auricomus* (Spach) Schur.

Morphologically, *Ranunculus jiguanishanicus* is somewhat similar to *R. pegaeus* (Figs 7–10), also a member of *R.* sect. *Ranunculus*, in having prostrate and glabrous stems (Figs 2A, B, 8A, B), small flowers (Figs 3D, 9D), subglobose aggregate




Figure 9. *Ranunculus pegaeus* in the wild (China, Sichuan, Maoxian) **A** roots **B** portion of stem **C** leaves **D** flower (lateral view) **E** flower (top view) **F** sepal (left: abaxial side; right: adaxial side) **G** petal (left: adaxial side; right: abaxial side) **H** stamens **I** carpels **J** aggregate fruit **K** achenes **L** receptacle. Photographed by Wen-Qun Fei.

中国科学院华南植物园标本采集记录

采集者: 费文群 采集编号: FWQ569
采集日期: 2022. 6. 7
采集地: 四川省茂县南新镇九鼎山
经纬度: N31° 30' 31.26" E103° 48' 6.57"
海拔高度: 3471 m 地形: 山地
生境: 杜鹃林下溪边苔藓丛中 分布: 少
习性: 多年生草本
根: 茎: 叶:
花: 黄色 果实/种子: 无毛
备注:
拉丁名: *Ranunculus pegaus* Hand.-Mazz.




Figure 10. *Ranunculus pegaus*. China, Sichuan, Maoxian, W.Q. Fei 569 (IBSC).

fruit (Figs 3I, 9I), and glabrous carpels (Figs 3H, 9H), achenes (Figs 3G, 9G) and receptacles (Figs 3K, 9K). However, it differs by having 3-foliolate leaves with obvious petiolules (3–5 mm long), unequally 3-sected leaflets, lanceolate to linear, entire or 1–2-denticulate to 1–2-cleft ultimate leaflet segments (Fig. 3C), and styles



Figure 11. Type sheet of *Ranunculus glareosus*.



Figure 12. *Ranunculus glareosus* in the wild (China, Qinghai, Menyuan) **A, B** habitat **C** habit. Photographed by Wen-Qun Fei.




Figure 13. *Ranunculus glareosus* in the wild (China, Qinghai, Menyuan) **A** roots **B** portion of stem **C** leaves **D** flower (lateral view) **E** flower (top view) **F** sepal (left: abaxial side; right: adaxial side) **G** petal (left: adaxial side; right: abaxial side) **H** stamens **I** carpels **J** aggregate fruit **K** achenes **L** receptacle. Photographed by Wen-Qun Fei.

in the carpels and achenes ca. 0.8 mm long (Fig. 3H, G). In *R. pegaus*, the leaves are 3-partite, 3-sected or 3-foliolate with the central segment/leaflet rhombic or oblong, entire or 3-denticulate and the lateral segments/leaflets obliquely flabellate, entire or unequally 2-cleft (Figs 9C, 10), and styles in the carpels and achenes ca. 0.3 mm long (Fig. 9H, G). A detailed morphological comparison between *R. jiguanishanicus* and *R. pegaus* is given in Table 1.

中国科学院华南植物园标本采集记录

采集者: 费文群 采集编号: FWQ623
采集日期: 2022. 6. 26
采集地: 青海省门源县达坂山
经纬度: N37° 20' 12.24" E101° 24' 8.97"
海拔高度: 4004 m 地形: 山地
生境: 流石滩 分布: 少
习性: 多年生草本
根: 茎: 叶:
花: 黄色 果实/种子: 无毛
备注:
拉丁名: *Ranunculus glareosus* Hand.-Mazz.



Ranunculus glareosus Hand.-Mazz.

鉴定人(Det.): 费文群 (Wen-qun Fei) 2022. 9. 11

Figure 14. *Ranunculus glareosus*. China, Qinghai, Menyuan, W.Q. Fei 623 (IBSC).

Table 1. Morphological comparison of *Ranunculus glareosus*, *R. jiguanshanicus* sp. nov. and *R. pegaus*.

	<i>R. glareosus</i>	<i>R. jiguanshanicus</i>	<i>R. pegaus</i>
Roots	2–5, more than 15 cm long	2–5, 6–10 cm long	5–10, 8–12 cm long
Stems	sparsely puberulous	glabrous	glabrous
Basal leaves	3-sected or 3-foliolate, fleshy, adaxially glabrous or sparsely puberulous, abaxially glabrous, central segment/leaflet ovate or rhombic, entire or 3-lobate, lateral segments/leaflets flabellate, unequally 2-partite	3-foliolate, thinly chartaceous, both sides glabrous, leaflets unequally 3-sected, with ultimate leaf segments narrowly lanceolate to linear, entire or 1–2-denticulate to 1–2-cleft	3-partite, 3-sected or 3-foliolate, thinly chartaceous, both sides glabrous, central segment/leaflet rhombic or oblong, entire or 3-denticulate, lateral segments/leaflets obliquely flabellate, entire or unequally 2-cleft
Flowers	terminal, 1–4, 15–17 mm in diameter	terminal, 1(–2), 5.2–6 mm in diameter	terminal or axillary, 3–7, 5.5–8 mm in diameter
Receptacles	3–5 mm long, clavate, glabrous	ca. 1.2 mm long, clavate, glabrous	ca. 1 mm long, clavate, glabrous
Sepals	adaxially glabrous, abaxially puberulous	both sides glabrous	both sides glabrous
Petals	9–10 × 7–8 mm, widely obovate	3.2–3.5 × 1.8–2 mm, obovate	3–3.5 × 1.5–1.7 mm, obovate
Carpels	20–35; ovaries ovoid, glabrous; styles ca. 0.2 mm long, straight	8–12; ovaries ovoid, glabrous; styles ca. 0.8 mm long, apex recurved	18–22; ovaries ovoid, glabrous; styles ca. 0.3 mm long, apex recurved
Aggregate fruit	ellipsoid	subglobose	subglobose
Achenes	widely ovoid, glabrous, styles ca. 0.2 mm long, straight	widely ovoid, glabrous, styles ca. 0.8 mm long, apex recurved	widely ovoid, glabrous, styles ca. 0.3 mm long, apex recurved

As mentioned earlier, a gathering of *Ranunculus jiguanshanicus*, W.B. Ju, L. Zhang & D.K. Chen AZH01290 (CDBI), from Chongzhou in Sichuan, the type locality of this species, had been previously misidentified as *R. glareosus* (Figs 11–14). Morphologically, *R. jiguanshanicus* is very easily distinguishable from *R. glareosus* by having glabrous stems (vs. sparsely puberulous) (Figs 3B, 13B), thinly chartaceous leaves (vs. fleshy), leaflets of the 3-foliolate leaves with obvious petiolules (3–5 mm vs. 0.5–2 mm long), unequally 3-sected, with the ultimate leaflet segments narrowly lanceolate to linear, entire or 1–2-denticulate to 1–2-cleft (vs. 3-sected or 3-foliolate, central segment/leaflet ovate or rhombic, entire or 3-lobed, and lateral segments/leaflets flabellate, unequally 2-partite) (Figs 3C, 13C), smaller flowers (5.2–6 mm vs. 15–17 mm in diameter) (Figs 3D, E, 13D, E), abaxially glabrous sepals (vs. puberulous) (Figs 3F, 13F), smaller (3.2–3.5 × 1.8–2 mm vs. 9–10 × 7–8 mm) and obovate petals (vs. widely obovate) (Figs 3G, 13G), subglobose aggregate fruit (vs. ellipsoid) (Figs 3J, 13J), and longer styles in the carpels and achenes (ca. 0.8 mm vs. ca. 0.2 mm long) (Figs 3I, K, 13I, K). In habitat, *R. jiguanshanicus* grows among moss on rocks or rocky cliffs in moist places in fir forests at altitudes of 2900–3150 m above sea level, whereas *R. glareosus* grows on alpine scree slopes at altitudes of 3900–4800 m above sea level. A detailed morphological comparison between *R. glareosus* and *R. jiguanshanicus* is given in Table 1.

Additional specimens examined (paratypes). CHINA. **Sichuan:** Chongzhou, W.B. Ju, L. Zhang & D.K. Chen AZH01290 (CDBI); Dayi, W.Q. Fei 897 (IBSC).

Acknowledgements

We are grateful to Dr. Andrey Erst, Dr. Andriy Novikov, and Dr. Marco Pellegrini, for their valuable comments on the manuscript. We thank the curators of CDBI, E, KUN, PE, and WU for allowing us to use their scanned images of specimens and for

research facilities. We also thank Jun Li and De-chang Meng for their assistance with the fieldwork. This work was supported by the National Natural Science Foundation of China (grant nos. 31870184, 31770218, 31970210).

References

- Fei WQ, Yuan Q, Yang QE (2022) *Ranunculus huainingensis* and *R. lujiangensis* (Ranunculaceae), described from Anhui in China, are both synonymous with *R. ternatus*, a polymorphic eastern Asian species. *Phytotaxa* 573(1): 15–38. <https://doi.org/10.11646/phytotaxa.573.1.2>
- Fei WQ, Yuan Q, Yang QE (2023a) *Ranunculus luanchuanensis* (Ranunculaceae), a new species from Henan, China. *PhytoKeys*. (in press)
- Fei WQ, Yuan Q, Yang QE (2023b) *Ranunculus maoxianensis* (Ranunculaceae), a new species from northwestern Sichuan, China, with an emended description of *R. chongzhouensis*, the putative closest ally of the new species. *PhytoKeys*. (in press)
- Handel-Mazzetti H (1931) Ranunculaceae. In: *Symbolae Sinicae* Vol. 7. Julius Springer, Wien, 265–321.
- Handel-Mazzetti H (1939) *Plantae sinensis a Dre. H. Smith annis 1921–1922, 1924 et 1934 lectae*. XXXIII. Ranunculaceae. *Acta Horti Gotoburgensis* 13: 37–219.
- Hörandl E, Emadzade K (2012) Evolutionary classification: A case study on the diverse plant genus *Ranunculus* L. (Ranunculaceae). *Perspectives in Plant Ecology, Evolution and Systematics* 14(2): 310–324. <https://doi.org/10.1016/j.ppees.2012.04.001>
- Hörandl E, Paun O, Johansson JT, Lehnbech C, Armstrong T, Chen L, Lockhart P (2005) Phylogenetic relationships and evolutionary traits in *Ranunculus* s.l. (Ranunculaceae) inferred from ITS sequence analysis. *Molecular Phylogenetics and Evolution* 36(2): 305–327. <https://doi.org/10.1016/j.ympev.2005.02.009>
- IUCN Standards and Petitions Committee (2022) Guidelines for Using the IUCN Red List Categories and Criteria. Version 14. IUCN Standards and Petitions Committee. <https://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Liou L (1980) *Ranunculus* L. In: Wang WT (Ed.) *Flora Reipublicae Popularis Sinicae* Vol. 28. Science Press, Beijing, 255–331.
- Luo MR, Zhao L (2013) A new *Ranunculus* species (Ranunculaceae) from Shaanxi, China. *Bangladesh Journal of Plant Taxonomy* 20(2): 201–205. <https://doi.org/10.3329/bjpt.v20i2.17394>
- Paun O, Lehnbech C, Johansson JT, Lockhart P, Hörandl E (2005) Phylogenetic relationships and biogeography of *Ranunculus* and allied genera (Ranunculaceae) in the Mediterranean region and in the European Alpine System. *Taxon* 54(4): 911–930. <https://doi.org/10.2307/25065478>
- Tamura M (1995) Angiospermae. Ordnung Ranunculales. Fam. Ranunculaceae. II. Systematic Part. In: Hiepko P (Ed.) *Natürliche Pflanzenfamilien*, 2nd edn. Vol. 17aIV. Duncker & Humblot, Berlin, 223–519.
- Thiers B (2022) Index Herbariorum: A Global Directory of Public Herbaria and Associated Staff. <http://sweetgum.nybg.org/science/ih/> [accessed 2 December 2022]

- Wang WT (1995a) A revision of the genus *Ranunculus* in China (I). *Bulletin of Botanical Research* 15(2): 137–180.
- Wang WT (1995b) A revision of the genus *Ranunculus* in China (II). *Bulletin of Botanical Research* 15(3): 275–329.
- Wang WT (1996) Notulae de Ranunculaceis Sinensibus (XIX). *Bulletin of Botanical Research* 16(2): 155–166.
- Wang WT (2007) *Ranunculus ailaoshanicus* W. T. Wang, a new species of Ranunculaceae from Yunnan, China. *Acta Phytotaxonomica Sinica* 45(3): 293–295. <https://doi.org/10.1360/aps06192>
- Wang WT (2008) New taxa of Ranunculaceae from Yunnan. *Acta Botanica Yunnanica* 30(5): 519–524.
- Wang WT (2013) Six new species and two new varieties of Ranunculaceae from southwest China. *Guihaia* 33(5): 579–587.
- Wang WT (2015) Five new species of *Ranunculus* from west China. *Bulletin of Botanical Research* 35(5): 641–646.
- Wang WT (2016) Six new species of Ranunculaceae from China. *Guihaia* 36(11): 1303–1311.
- Wang WT (2018) *Ranunculus lujiangensis*, a new species of Ranunculaceae from Anhui Province. *Bulletin of Botanical Research* 38(6): 801–803. <https://doi.org/10.7525/j.issn.1673-5102.2018.06.001>
- Wang WT (2019a) *Ranunculus kangmaensis* W. T. Wang, a new species of Ranunculaceae from Tibet of China. *Guihaia* 39(3): 285–287. <https://doi.org/10.11931/guihaia.gxzw201712013>
- Wang WT (2019b) Two new species of *Ranunculus* from Tibet, China. *Guihaia* 39(9): 1139–1342. <https://doi.org/10.11931/guihaia.gxzw201811047>
- Wang WT (2022) Five new species and one new variety of *Ranunculus* (Ranunculaceae) from Sichuan, with one new section represented by one of these species. *Guihaia* 42(1): 1–9. <https://doi.org/10.11931/guihaia.gxzw202011047>
- Wang WT, Chen SR (2015) *Ranunculus laohegouensis*, a new species of Ranunculaceae from Sichuan. *Bulletin of Botanical Research* 38(6): 801–802. <https://doi.org/10.7525/j.issn.1673-5102.2015.06.001>
- Wang WT, Gilbert MG (2001) *Ranunculus* Linnaeus. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* Vol. 6. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 391–431.
- Wang WT, Liao L (2009) *Ranunculus napahaiensis*, a new species of Ranunculaceae from Yunnan, China. *Guihaia* 29(4): 427–429.
- Wang WT, Yang Z, Xie J (2016) *Ranunculus huainingensis*, a new species of Ranunculaceae from Anhui. *Guihaia* 36(supplement 1): 97–99.
- Yang QE (2000) *Ranunculus wangianus* Q. E. Yang, a new species from NW Yunnan, China and its karyotype. *Acta Phytotaxonomica Sinica* 38(6): 551–556.
- Yuan Q, Yang QE (2017a) *Ranunculus angustisepalus* (Ranunculaceae) is an *Oxygraphis* and conspecific with *O. delavayi*. *Phytotaxa* 319(1): 103–110. <https://doi.org/10.11646/phytotaxa.319.1.6>

- Yuan Q, Yang QE (2017b) The identity of *Ranunculus ailaoshanicus* (Ranunculaceae) from China. *Phytotaxa* 319(1): 111–117. <https://doi.org/10.11646/phytotaxa.319.1.7>
- Yuan Q, Yang QE (2017c) The identity of *Ranunculus laohegouensis* (Ranunculaceae) from Sichuan, China. *Phytotaxa* 324(2): 198–200. <https://doi.org/10.11646/phytotaxa.324.2.9>
- Zhang M, Zeng YP, Yang QE (2020) *Ranunculus kangmaensis* (Ranunculaceae), a new synonym of *Halerpestes tricuspis* var. *variifolia*. *Phytotaxa* 434(1): 101–112. <https://doi.org/10.11646/phytotaxa.434.1.7>